

QCD ASPECTS OF CDF RESULTS ON DIFFRACTION

QCD 2004

7-11 June 2004

L'Institut d'Astrophysique de Paris
Paris, France

Konstantin Goulianos

The Rockefeller University


- CDF results
- Comparison with HERA
- QCD aspects




Forty Years of Diffraction

- ✚ 1960's Good and Walker
BNL: first observation
- ✚ 1970's Fermilab fixed target, ISR, SPS
Regge factorization works
KG, Phys. Rep. 101, 169 (1983)
- 1980's UA8: diff. dijets \Rightarrow hard diffraction
- 1990's Tevatron: Regge factorization breakdown
Tev, HERA: QCD factorization breakdown

Pre-CDF Soft Diffraction



QCD Aspects




$$\xi = \frac{\Delta p_L}{p_L} = \frac{M^2}{s}$$

$$\frac{d\sigma}{dM^2} \sim \frac{1}{M^2} \Leftrightarrow \frac{d\sigma}{d\Delta y} \propto \text{constant}$$

POMERON: color singlet w/vacuum quantum numbers

$$\left. \frac{d^2\sigma}{dt dM^2} \right|_{t=0.025}$$




KG, Phys. Rep. 101 (1983) 171

Diffraction@CDF in Run I

16 papers


- Elastic scattering PRD 50 (1994) 5518
- Total cross section PRD 50 (1994) 5550
- Diffraction

SOFT diffraction



HARD diffraction

PRL references



with roman pots

JJ 84 (2000) 5043
JJ 88 (2002) 151802

W 78 (1997) 2698	JJ 74 (1995) 855	JJ 85 (2000) 4217
JJ 79 (1997) 2636	JJ 80 (1998) 1156	
b-quark 84 (2000) 232	JJ 81 (1998) 5278	
J/ψ 87 (2001) 241802		


Total & Elastic Cross Sections

(Run I-0)

Total and Elastic Cross Sections

Covolan, Montanha and Goulianos, Phys. Lett. B 389 (1996) 176

$$\alpha_{pp} = 1 + \epsilon (\Rightarrow 0.104) + 0.25t \quad \alpha_{p\bar{p}} = 0.68 + 0.82t \quad \alpha_{pp} = 0.46 + 0.92t$$




$$\sigma_T(s) = \sigma_0 s^\epsilon = \sigma_0 e^{\epsilon \Delta y'}$$

The exponential rise of σ_T is a QCD aspect expected in the parton model
(see E. Levin, An Introduction to Pomerons, Preprint DESY 98-120)

$$\text{Im } f_{el}(s, t) \propto e^{(\epsilon + \alpha' t) \Delta y}$$

Soft Diffraction (Run I-0)



$$\frac{d^2\sigma_{SD}}{dt d\xi} = f_{IP/p}(t, \xi) \cdot \sigma_{IP-\bar{p}}(M_X^2)$$

$$\sigma_{SD} \sim S^{2\varepsilon}$$

❖ Unitarity problem:


With factorization and std pomeron flux σ_{SD} exceeds σ_T at $\sqrt{s} \approx 2 \text{ TeV}$.

❖ Renormalization:

normalize the pomeron flux to unity

KG, PLB 358 (1995) 379

$$\int_{\xi_{\min}}^{0.1} \int_{t=-\infty}^0 f_{IP/p}(t, \xi) d\xi dt = 1$$




A Scaling Law in Diffraction

Factorization breaks down in favor of M^2 -scaling


renormalization

$$\frac{d\sigma}{dM^2} \propto \frac{s^{2\varepsilon} \rightarrow 1}{(M^2)^{1+\varepsilon}}$$

KG&JM, PRD 59 (1999) 114017




Central and Double Gaps




□ Double Diffraction Dissociation

➤ One central gap



□ Double Pomeron Exchange

➤ Two forward gaps




□ SDD: Single+Double Diffraction

➤ One forward + one central gap

QCD Basis of Renormalization

(KG, hep-ph/0205141)



2 independent variables: $t, \Delta y$

color factor

$$\kappa = \frac{g_{IP-IP-IP}(t)}{\beta_{IP-p-p}(0)} \approx 0.17$$

$$\frac{d^2\sigma}{dt d\Delta y} = \underbrace{C \cdot F_p^2(t_1)}_{\text{Gap probability}} \cdot \left\{ e^{(\varepsilon + \alpha' t)\Delta y} \right\}^2 \cdot \underbrace{\kappa \cdot \left\{ \sigma_0 e^{\varepsilon \Delta y'} \right\}}_{\text{color factor}}$$

Gap probability

$$\sim e^{2\varepsilon \Delta y}$$

$$\int_{\Delta y_{\min}}^{\Delta y = \ln s} s^{2\varepsilon \Delta y} \approx s^{2\varepsilon}$$

Renormalization removes the s-dependence → SCALING

The Factors K and ε


Experimentally:

$$K = \frac{g_{IP-IP-IP}}{\beta_{IP-p}} = 0.17 \pm 0.02, \quad \varepsilon = 0.104$$

KG&JM, PRD 59 (114017) 1999


Color factor: $K = f_g \times \frac{1}{N_c^2 - 1} + f_q \times \frac{1}{N_c} \xrightarrow{Q^2=1} \approx 0.75 \times \frac{1}{8} + 0.25 \times \frac{1}{3} = 0.18$

Pomeron intercept: $\varepsilon = \lambda_g \cdot w_g + \lambda_q \cdot w_q = 0.12$




$$X \cdot f(X) = \frac{1}{X^\lambda}$$

f_g = gluon fraction
 f_q = quark fraction




Generalized Renormalization

(KG, hep-ph/0205141)



5 independent variables



y'_1 y_2

t_1 $\Delta y = \Delta y_1 + \Delta y_2$ t_2

color factors

$$\frac{d^5 \sigma}{\prod_{i=1-5} dV_i} = C \times F_p^2(t_1) \prod_{i=1-2} \left\{ e^{(\varepsilon + \alpha' t_i) \Delta y_i} \right\}^2 \times \kappa^2 \left\{ \sigma_o e^{\varepsilon(\Delta y'_1 + \Delta y'_2)} \right\}$$

Gap probability
 $\sim e^{2\varepsilon \Delta y}$


Sub-energy cross section
 (for regions with particles)

$$\int_{\Delta y_{\min}}^{\Delta y = \ln s} s^{2\varepsilon \Delta y} \approx s^{2\varepsilon}$$


Same suppression as for single gap!

Central & Double-Gap Results


Differential shapes agree with Regge predictions



- One-gap cross sections are suppressed
- Two-gap/one-gap ratios are $\approx \kappa = 0.17$



Soft Gap Survival Probability



$$S = \frac{\phi \left[\begin{array}{|c|c|c|} \hline \eta & & \eta \\ \hline \end{array} \right] / \phi \left[\begin{array}{|c|} \hline \eta \\ \hline \end{array} \right]}{\phi \left[\begin{array}{|c|c|c|} \hline \eta & & \eta \\ \hline \end{array} \right] / \phi \left[\begin{array}{|c|c|c|} \hline \eta & & \eta \\ \hline \end{array} \right]}$$

$$S_{2\text{-gap}/1\text{-gap}}^{1\text{-gap}/0\text{-gap}} (1800 \text{ GeV}) \approx 0.23$$

$$S_{2\text{-gap}/1\text{-gap}}^{1\text{-gap}/0\text{-gap}} (630 \text{ GeV}) \approx 0.29$$

Soft Diffraction Conclusions

Experiment:



- M^2 - scaling
- Non-suppressed double-gap to single-gap ratios

Phenomenology:

- Generalized renormalization
- Obtain Pomeron intercept and tripe-Pomeron coupling from inclusive PDF's and color factors

Soft and Hard Diffraction

SOFT DIFFRACTION





$$\xi = \Delta P_L / P_L$$

ξ = fractional momentum loss
of scattered (anti)proton

Variables: (ξ, t) or $(\Delta\eta, t)$

HARD DIFFRACTION



Additional variables: (x, Q^2)



$$x_{Bj} = \sum E_T^{jet} e^{-\eta^{jet}} / \sqrt{s}$$

$$x = \beta \xi, \quad Q^2 = (E_T^{jet})^2$$

Run-IC

CDF-I

Run-IA,B



Forward Detectors

BBC $3.2 < \eta < 5.9$

FCAL $2.4 < \eta < 4.2$

Diffraction Fractions




$$\bar{p}p \rightarrow X + \text{gap}$$

SD/ND fraction at 1800 GeV

X	Fraction(%)
W	1.15 (0.55)
JJ	0.75 (0.10)
b	0.62 (0.25)
J/ψ	1.45 (0.25)

$$\bar{p}p \rightarrow \text{Jet} + \text{gap} + \text{Jet}$$

DD/ND gap fraction at 1800 GeV



- All SD/ND fractions ~1%
- Gluon fraction $f_g = 0.54 \pm 0.15$
- Suppression by ~5 relative to HERA
→ gap survival probability ~20%

Factorization OK @ Tevatron
at 1800 GeV (single energy) ?

Diffractive Structure F'n @CDF

$$\bar{p} + p \rightarrow \bar{p} + Jet + Jet + X$$

- Measure ratio of SD/ND dijet rates as a f'n of $x_{\bar{p}}$


$$x_{\bar{p}} \equiv p_{g,q}/p_{\bar{p}} = \frac{\sum_{i=1}^{2(3)} E_T^i \cdot e^{-\eta^i}}{\sqrt{s}}$$

$$\frac{R_{SD}}{R_{ND}}(x_{\bar{p}}) \approx R_0 \cdot x_{\bar{p}}^{-0.45}$$

- In LO-QCD ratio of rates equals ratio of structure f'n's




$$F_{jj}(x_{\bar{p}}) = x_{\bar{p}} \left[g(x_{\bar{p}}) + \frac{C_F}{C_A} \sum (q_i(x_{\bar{p}}) + \bar{q}_i(x_{\bar{p}})) \right]$$

SD/ND Rates vs $x_{\bar{p}}$



Breakdown of QCD Factorization

HERA $\xrightarrow{\text{The clue to understanding the Pomeron}}$ TEVATRON



$$F_2(Q^2, x)$$



$$F_{JJ}(E_T^{\text{Jet}}, x)$$

$$F_2^D(Q^2, \beta, \xi, t)$$


???

$$F_{JJ}^D(E_T^{\text{Jet}}, \beta, \xi, t)$$


Restoring Diffractive Factorization




DSF from single-gaps




DSF from double-gaps:
Factorization restored!



Q² dependence of DSF




- $F_2^{D(3)}(x_{IP}, \beta, Q^2)$ H1 prel.
- $F_2(x, Q^2)$ H1 96-97



$$R \left(\frac{F^D(Q^2, x, \xi)}{F(Q^2, x)} \right) \Rightarrow \begin{cases} \sim \text{no } Q^2 \text{ dependence} \\ \sim \text{flat at HERA} \\ \sim 1/x^{0.5} \text{ at Tevatron} \end{cases}$$

Pomeron evolves similarly to proton
except for for renormalization effects

Diffractive Structure Function from Inclusive pdf's (KG)




$$x \cdot f(x) = \frac{1}{x^\epsilon}$$

Power-law region

$$\xi_{\max} = 0.1$$

$$x_{\max} = 0.1$$

$$\beta < 0.05\xi$$



$$F^D(\varrho^2, x, \xi) \propto \frac{1}{\xi^{1+\epsilon}} \cdot F(\varrho^2, x) \propto \frac{1}{\xi^{1+\epsilon}} \cdot \frac{C(\varrho^2)}{(\beta\xi)^{\lambda(\varrho^2)}} \Rightarrow \frac{A_{\text{NORM}}}{\xi^{1+\epsilon+\lambda}} \cdot \kappa \cdot \frac{C}{\beta^\lambda}$$


HERA(no RENORM): $R_{DIS}^{DDIS}(x) \xrightarrow{\text{fixed } \xi} \text{constant}$

TEVATRON (RENORM) : $R_{ND}^{SD}(x) \propto x^{-(\epsilon + \lambda)}$



$$2\epsilon_{DDIS} = \epsilon + \lambda(Q^2)$$

Pomeron Intercept from H1

H1 Diffractive Effective $\alpha_{IP}(0)$ $\alpha_{IP}(t) = 1 + \varepsilon + \alpha' t$



ξ -dependence: Inclusive vs Dijets



$$\frac{d\sigma_{\text{incl}}}{d\xi} \propto \text{constant}$$


$$F_{jj}^D(\beta, \xi) \propto \frac{1}{\beta^n} \cdot \frac{1}{\xi^m} \quad (n = 1.0 \pm 0.1, \quad m = 0.9 \pm 0.1)$$

Pomeron dominated

Exclusive Dijets in DPE

Interest in diffractive Higgs production

Calibrate on exclusive dijets




Dijet mass fraction

$$R_{jj} = \frac{M_{jj}^{\text{cone}}}{M_X}$$

E_T^{jet}	$\sigma_{\text{DPE}}^{\text{excl jj}} (R_{jj} > 0.8)$
10 GeV	$970 \pm 65 \pm 272 \text{ pb}$
25 GeV	$34 \pm 5 \pm 10 \text{ pb}$

Upper limit for excl DPE-jj consistent with theory: KMR $\rightarrow 60 \text{ pb} @ 25 < E_T^J < 35 \text{ GeV}$



CDF Run II Preliminary



Search for Exclusive χ_c @CDF

$$\bar{p} + p \rightarrow \bar{p} + \chi_c (\rightarrow J/\psi + \gamma) + p$$






- Events are triggered on dimuons
- Select muons with $P_T > 1.5 \text{ GeV}$, $|\eta| < 0.6$
- Reject cosmic rays using time of flight information
- Select events in J/ψ mass window



No positive identification of χ_c events
Cross section upper limit comparable to KMR prediction

Gap Between Jets

$\bar{p} + p \rightarrow \text{Jet} + \text{Gap} + \text{Jet}$



Summary

SOFT DIFFRACTION

- M^2 - scaling
- Non-suppressed double-gap to single-gap ratios

HARD DIFFRACTION

- Flavor-independent SD/ND ratio
- Little or no Q^2 -dependence in SD/ND ratio

- ✓ Universality of gap prob. across soft and hard diffraction
- ✓ Pomeron evolves similarly to proton

Diffraction appears to be a low- x partonic exchange subject to color constraints