

Diffraction cross sections implemented in PYTHIA8-MBR vs LHC results

KONSTANTIN GOULIANOS

The Rockefeller University, 1230 York Avenue, New York, NY 10065, USA

Diffraction cross sections implemented in the PYTHIA8-MBR Monte Carlo simulation are reviewed and compared to results obtained at the Large Hadron Collider.

1. Introduction

Measurements at the LHC have shown that there are sizable disagreements among Monte Carlo (MC) implementations of “soft” processes based on cross sections proposed by various physics models, and that it is not possible to reliably predict all such processes, or even all aspects of a given process, using a single model [1, 2, 3]. In the CDF studies of diffraction at the Tevatron, all processes are well modeled by the MBR (Minimum Bias Rockefeller) MC simulation, which is a stand-alone simulation based on a unitarized Regge-theory model, RENORM [4], employing inclusive nucleon parton distribution functions (PDF’s) and QCD color factors. The RENORM model was updated in a presentation at EDS-2009 [5] to include a unique unitarization prescription for predicting the total pp cross section at high energies, and that update has been included as an MBR option for simulating diffractive processes in PYTHIA8 since version PYTHIA8.165 [6], to be referred here-forth as PYTHIA8-MBR. In this paper, we briefly review the cross sections [7] implemented in this option of PYTHIA8 and compare them with LHC measurements.

The PYTHIA8-MBR option includes a full simulation of the hadronization of the implemented diffraction dissociation processes: single, double, and central diffraction. In the original MBR simulation used in CDF, the hadronization of the final state(s) was based on a data-driven phenomenological model of multiplicities and p_t distributions calibrated using $S\bar{p}pS$ and Fermilab fixed-target results. Later, the model was successfully tested against Tevatron MB and diffraction data. However, only π^\pm and π^0 particles were produced in the final state, with multiplicities obeying a statistical model of a modified Gamma distribution function that provided good fits

to experimental data [8]. This model could not be used to predict specific-particle final states. In the PYTHIA8-MBR implementation, hadronization is performed by PYTHIA8 tuned to reproduce final-state distributions in agreement with MBR's, with hadronization done in the PYTHIA8 framework. Thus, all final-state particles are now automatically produced, greatly enhancing the horizon of applicability of tPYTHIA8-MBR.

2. Cross sections

The following diffraction dissociation processes are considered in PYTHIA8-MBR:

$$\text{SD } pp \rightarrow Xp \quad \text{Single Diffraction (or Single Dissociation),} \quad (1)$$

$$\text{or } pp \rightarrow pY \quad (\text{the other proton survives})$$

$$\text{DD } pp \rightarrow XY \quad \text{Double Diffraction (or Double Dissociation),} \quad (2)$$

$$\text{CD/DPE } pp \rightarrow pXp \quad \text{Central Diffraction/Double Pomeron Exchange.} \quad (3)$$

The RENORM predictions are expressed as unitarized Regge-theory formulas, in which the unitarization is achieved by a renormalization scheme where the Pomeron (\mathbb{P}) flux is interpreted as the probability for forming a diffractive (non-exponentially suppressed) rapidity gap and thereby its integral over all phase space saturates at the energy where it reaches unity. Differential cross sections are expressed in terms of the \mathbb{P} -trajectory, $\alpha(t) = 1 + \epsilon + \alpha' t = 1.104 + 0.25 \text{ (GeV}^{-2}) \cdot t$, the \mathbb{P} - p coupling, $\beta(t)$, and the ratio of the triple- \mathbb{P} to the \mathbb{P} - p couplings, $\kappa \equiv g(t)/\beta(0)$. For large rapidity gaps, $\Delta y \geq 3$, for which \mathbb{P} -exchange dominates, the cross sections may be written as,

$$\frac{d^2\sigma_{SD}}{dt d\Delta y} = \frac{1}{N_{\text{gap}}(s)} \left[\frac{\beta^2(t)}{16\pi} e^{2[\alpha(t)-1]\Delta y} \right] \cdot \left\{ \kappa \beta^2(0) \left(\frac{s'}{s_0} \right)^\epsilon \right\}, \quad (4)$$

$$\frac{d^3\sigma_{DD}}{dt d\Delta y dy_0} = \frac{1}{N_{\text{gap}}(s)} \left[\frac{\kappa \beta^2(0)}{16\pi} e^{2[\alpha(t)-1]\Delta y} \right] \cdot \left\{ \kappa \beta^2(0) \left(\frac{s'}{s_0} \right)^\epsilon \right\}, \quad (5)$$

$$\frac{d^4\sigma_{DPE}}{dt_1 dt_2 d\Delta y dy_c} = \frac{1}{N_{\text{gap}}(s)} \left[\prod_i \frac{\beta^2(t_i)}{16\pi} e^{2[\alpha(t_i)-1]\Delta y_i} \right] \cdot \kappa \left\{ \kappa \beta^2(0) \left(\frac{s'}{s_0} \right)^\epsilon \right\}, \quad (6)$$

where t is the 4-momentum-transfer squared at the proton vertex, Δy the rapidity-gap width, and y_0 the center of the rapidity gap. In Eq. (6), the subscript $i = 1, 2$ enumerates Pomerons in a DPE event, $\Delta y = \Delta y_1 + \Delta y_2$ is the total rapidity gap (sum of two gaps) in the event, and y_c is the center in η of the centrally-produced hadronic system.

The total cross section (σ_{tot}) is expressed as:

$$\sigma_{\text{tot}}^{p\pm p} = 16.79s^{0.104} + 60.81s^{-0.32} \mp 31.68s^{-0.54} \quad \text{for } \sqrt{s} \leq 1.8 \text{ TeV}, \quad (7)$$

$$\sigma_{\text{tot}}^{p\pm p} = \sigma_{\text{tot}}^{\text{CDF}} + \frac{\pi}{s_0} \left[\left(\ln \frac{s}{s_F} \right)^2 - \left(\ln \frac{s^{\text{CDF}}}{s_F} \right)^2 \right] \quad \text{for } \sqrt{s} \geq 1.8 \text{ TeV}, \quad (8)$$

where s_0 and s_F are energy and the Pomeron flux saturation scales, respectively [7]. For $\sqrt{s} \leq 1.8$ TeV, where there are Reggeon contributions, we use the global fit expression [9], while for $\sqrt{s} \geq 1.8$ TeV, where Reggeon contributions are negligible, we employ the Froissart-Martin formula [10, 11, 12]. The two expressions are smoothly matched at $\sqrt{s} \sim 1.8$ TeV.

The elastic cross section is obtained from the global fit [9] for $\sqrt{s} \leq 1.8$ TeV, while for $1.8 < \sqrt{s} \leq 50$ TeV we use an extrapolation of the global-fit ratio of $\sigma_{\text{el}}/\sigma_{\text{tot}}$, which is slowly varying with \sqrt{s} , multiplied by σ_{tot} . The total non-diffractive cross section is then calculated as $\sigma_{\text{ND}} = (\sigma_{\text{tot}} - \sigma_{\text{el}}) - (2\sigma_{\text{SD}} + \sigma_{\text{DD}} + \sigma_{\text{CD}})$.

3. Results

In this section, we present as examples of the predictive power of the RENORM model some results reported by the TOTEM, CMS, and ALICE collaborations for pp collisions at $\sqrt{s} = 7$ TeV, which can be directly compared with RENORM formulas without using the PYTHIA8-MBR simulation. Figure 1 (top) shows a comparison of the TOTEM total, elastic, and total-inelastic cross sections, along with results from other experiments fitted by the COMPETE Collaboration [13]; the RENORM predictions, displayed as filled (green) squares, are in excellent agreement with the TOTEM results. Similarly, in Fig. 1 (bottom), good agreement is observed between the ALICE [14] and CMS [15] total-inelastic cross sections and the RENORM prediction.

The uncertainty shown in the RENORM prediction of σ_{tot} in Fig. 1 (top) is dominated by that in the scale parameter s_0 . The latter can be reduced by a factor of ~ 4 if $\sqrt{s_0}$ is interpreted as the mean value of the glue-ball-like object discussed in [16] and the data shown in Fig. 8 of [16] are used to determine its value. Work is in progress to finalize the details of this interpretation.

Another example of the predictive power of RENORM is shown in Fig. 2, which displays the total SD (top) and total DD (bottom) cross sections for $\xi < 0.05$, after extrapolation into the low mass region from the measured CMS cross sections at higher mass regions, presented in [17], using RENORM.

The justification for using the RENORM model for the extrapolation into the low mass region is presented in Fig. 3, in which the measured diffractive cross sections within a wide (albeit limited) pseudorpidity regions are






Fig. 1. (top) TOTEM measurements of the total, total-inelastic, and elastic pp cross sections at $\sqrt{s} = 7$ TeV shown with best COMPETE fits [13] and RENORM predictions; (bottom) ALICE [14] and CMS [15] measurements of the total inelastic cross section at $\sqrt{s} = 7$ TeV compared to RENORM prediction (PYTHIA8-MBR).



KG*: the value of this entry was obtained by extrapolating from the measured CMS cross sections [17] to the low mass region using the MBR model.

Fig. 2. Measured SD (top) and DD (bottom) cross sections for $\xi < 0.05$ compared with theoretical predictions; the model embedded in PYTHIA8-MBR provides a good description of all data.

compared with predictions. In Figs. 3 (top-left and top-middle), the predictions of PYTHIA8-MBR are shown for two values of the ϵ parameter of the Pomeron trajectory ($\alpha(t) = 1 + \epsilon + \alpha' t$), $\epsilon = 0.08$ and $\epsilon = 0.104$. Both values describe the measured SD cross section within uncertainties, while the DD data favor the smaller value of ϵ , which is consistent with low mass CDF data. The predictions of PYTHIA8-4C and PYTHIA6 describe well the measured DD cross section, but fail to describe the falling behavior of the data (see details in [17]). The total SD cross section integrated over the region $-5.5 < \log_{10} \xi < -2.5$ ($12 < M_X < 394$ GeV) was measured to be $\sigma_{vis}^{SD} = 4.27 \pm 0.04(\text{stat.})_{-0.58}^{+0.65}(\text{syst.})$ mb (dissociation of either proton).

The event sample after the $\Delta\eta^0 > 3$ selection, was used to extract the differential DD cross section as a function of the central-gap width, $\Delta\eta$. The cross section for $\Delta\eta > 3$, $M_X > 10$ GeV and $M_Y > 10$ GeV is presented in Fig. 3 (bottom). The total DD cross section integrated over this region was measured to be $\sigma_{vis}^{DD} = 0.93 \pm 0.01(\text{stat.})_{-0.22}^{+0.26}(\text{syst.})$ mb.




Fig. 3. SD (top) and DD (middle) cross sections vs ξ , and DD cross section vs $\Delta\eta$ (bottom), compared to PYTHIA6, PYTHIA8-4C and PYTHIA8-MBR. Errors are dominated by systematics (HF energy scale, and hadronization/diffraction model).

4. Summary

We reviewed our pre-LHC predictions for the total, elastic, total-inelastic, and diffractive components of the proton-proton cross section at high energies, which are based on a special parton-model approach to diffraction employing inclusive proton parton distribution functions and QCD color factors. We discuss single diffraction/dissociation, double diffraction/dissociation, and central diffraction or double-Pomeron exchange, comparing predictions with LHC measurements. Agreement is found in all cases.

Acknowledgments

I would like to thank Robert Ciesielski, my colleague at Rockefeller and collaborator in the implementation of the MBR simulation into PYTHIA8, and the Office of Science of the Department of Energy for supporting the Rockefeller experimental diffraction physics programs at Fermilab and LHC on which this research is anchored.

References

- [1] K. Goulianos, *Predictions of Diffractive Cross Sections in Proton-Proton Collisions*, in proceedings of *Diffraction 2012: International Workshop on Diffraction in High Energy Physics, 10-15 September 2012*, AIP Conf. Proc. 1523, pp. 107-110 (2013) [doi:<http://dx.doi.org/10.1063/1.4802128>].
- [2] See models presented by various authors in proceedings of *Diffraction 2012*, AIP Conf. Proc. **1523**.
- [3] K. Goulianos, *Predictions of Diffractive, Elastic, Total, and Total-Inelastic pp Cross Sections vs LHC Measurements*, to appear in proceedings of *XXI International Workshop on Deep-Inelastic Scattering and Related Subject - DIS2013, 22-26 April 2013, Marseilles, France*.
- [4] K. Goulianos, *Hadronic Diffraction: Where do we Stand?*, in proceedings of *Les Rencontres de Physique de la Vallée d'Aoste: Results and Perspectives in Particle Physics, La Thuile, Italy, February 27 - March 6, 2004*, Frascati Physics Series, Special 34 Issue, edited by Mario Greco [arXiv:[hep-ph/0407035](http://arxiv.org/abs/hep-ph/0407035)].
- [5] K. Goulianos, *Diffractive and Total pp Cross Sections at LHC*, in proceedings of *13th International Conference on Elastic and Diffractive Scattering (Blois Workshop) - Moving Forward into the LHC Era, CERN, Geneva, Switzerland, June 29-July 3, 2009*, CERN-Proceedings-2010-02, edited by Mario Deile [arXiv:[1002.3527v2](http://arxiv.org/abs/1002.3527v2)].
- [6] T. Sjöstrand, S. Mrenna and P. Skands, *JHEP05 (2006) 026, Comput. Phys. Comm. 178 (2008) 852* [arXiv:[hep-ph/0603175](http://arxiv.org/abs/hep-ph/0603175), arXiv:[0710.3820](http://arxiv.org/abs/0710.3820)].
- [7] R. Ciesielski and K. Goulianos, *MBR Monte Carlo Simulation in PYTHIA8* [arXiv:[1205.1446](http://arxiv.org/abs/1205.1446)].

- [8] K. Goulianos, Phys. Lett. B **193**, 151 (1987).
- [9] R. J. M. Covolan, J. Montanha and K. Goulianos, Phys. Lett. B **389**, 176 (1996).
- [10] M. Froissart, Phys. Rev. **3**, 123 (1961).
- [11] A. Martin, Nuovo Cimento **42**, 930 (1966).
- [12] A. Martin, Phys. Rev. D **80**, 065013 (2009).
- [13] M. Deile (TOTEM Collaboration), *Measurements of Proton-Proton and Total Cross Section at the LHC by TOTEM*, in proceedings of DIFFRACTION 2012 (figure from talk presented at the Conference).
- [14] B. Abelev et al. (ALICE Collaboration), *Measurement of Inelastic, Single- and Double-Diffraction Cross Sections in Proton-Proton Collisions at the LHC with ALICE*, Eur. Phys. J. C (2013)73:2456, [arXiv:1208.4968], [DOI 10.1140/cpjc/s10052-013-2456-0]
- [15] S. Chatrchyan et al. (CMS Collaboration), *Measurement of the inelastic proton-proton cross section at $\sqrt{s} = 7$ TeV*, Phys. Lett. B **722**, 5 (2013).
- [16] M. G. Albrow, T. D. Coughlin, J. R. Forshaw, *Central Exclusive Particle Production at High Energy Hadron Colliders*, Prog. Part. Nucl. Phys **65**, 149-184, 2010 [DOI: 10.1016/j.pnpnp.2010.06.001].
- [17] R. Ciesielski (CMS Collaboration), *Measurements of diffraction in p-p collisions in CMS*, in proceedings of XXI International Workshop on Deep-Inelastic Scattering and Related Subject -DIS2013, 22-26 April 2013, Marseilles, France.